



# Elliptic Sombor Index of Graphs From Primary Subgraphs

Nima Ghanbari<sup>1,\*</sup>, Saeid Alikhani<sup>1</sup>

<sup>1</sup> Department of Mathematical Sciences, Yazd University, 89195-741, Yazd, Iran

\* Corresponding author(s): [alikhani@yazd.ac.ir](mailto:alikhani@yazd.ac.ir)

Received: 21/06/2024 Revised: 30/07/2024 Accepted: 30/07/2024 Published: 02/08/2024



10.22128/gadm.2024.842.1115

## Abstract

Suppose that  $G$  is a connected graph constructed from pairwise disjoint connected graphs  $G_1, \dots, G_t$  by selecting a vertex of  $G_1$ , a vertex of  $G_2$ , and identifying these two vertices. Then continue in this manner inductively. The graphs  $G_1, \dots, G_k$  are the primary subgraphs of  $G$ . Some particular cases of these graphs are important in chemistry which we consider them in this paper and study their elliptic Sombor index.

**Keywords:** Sombor index, Elliptic Sombor index, Graph, Polymer

**Mathematics Subject Classification (2020):** 05C31

## 1 Introduction

A molecular graph is a simple graph such that its vertices correspond to the atoms and the edges to the bonds of a molecule. Suppose that  $G = (V, E)$  is a finite, connected, simple graph. As usual the degree of a vertex  $v$  in  $G$  is denoted by  $d_v$ .

The topological indices are the numerical parameters associated with the graph which are usually graph invariant. The topological index of a graph is based on the properties of graphs such as degree, distance, number of non-incident edges and so on. From this index it is possible to analyze the mathematical values and further investigate some physicochemical properties of a molecule. Therefore, it is also called a molecular descriptor. The first distance based topological index, is Wiener index


$$W(G) = \sum_{\{u,v\} \subseteq G} d(u,v) = \frac{1}{2} \sum_{u,v \in V(G)} d(u,v),$$

with the summation runs over all pairs of vertices of  $G$  [27]. The Wiener index is one of the most used topological indices with high correlation with many physical and chemical indices of molecular compounds [27]. The Sombor index which is a vertex-degree-based molecular structure descriptor introduced by Gutman in [16] and is defined as

$$SO(G) = \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2}.$$

In a remarkably brief period, the Sombor index has garnered considerable attention from both mathematicians and theoretical chemists. Redžepović [23] delved into its efficacy in prognosticating alkanes' entropy as well as enthalpy of vaporization, utilizing statistical analyzing





**Figure 1.** A graph  $G$  obtained by point-attaching from  $G_1, \dots, G_t$ .

techniques. Owing to its notably enhanced predictive capabilities, the Sombor index is adopted regarding the purpose of modeling thermodynamic properties of organic molecular structures [19]. For more details and aspects on the Sombor index we refer the reader to [1, 5, 6, 8, 12, 14, 15, 23, 26].

In [17] a novel geometric method is proposed for constructing vertex-degree-based molecular structure descriptors (topological indices). The model is based on an ellipse whose focal points represent the degrees of a pair of adjacent vertices. The approach enables a geometric interpretation of several previously known topological indices, and lead to design of a few new. The area of the ellipse induces a vertex-degree-based topological index of remarkable simplicity, which is called elliptic Sombor index. In [17]), the elliptic Sombor index (ESO) of  $G$  is defined as

$$ESO(G) = \sum_{uv \in E(G)} (d_u + d_v) \sqrt{d_u^2 + d_v^2}.$$

In [10], the extremal value problem for ESO over the set of (connected) graphs with equal number of vertices has studied. Also, the elliptic Sombor energy has investigated in [2].

Suppose that  $G$  is a connected graph constructed from pairwise disjoint connected graphs  $G_1, \dots, G_t$  as follows. Select a vertex of  $G_1$ , a vertex of  $G_2$ , and identify these two vertices. Then continue in this manner inductively. Note that the graph  $G$  constructed in this way has a tree-like structure, the  $G_i$ 's being its building stones (see Figure 1). The graphs  $G_1, \dots, G_k$  are the primary subgraphs of  $G$ . Usually say that  $G$  is a graph (polymer graph), obtained by point-attaching from  $G_1, \dots, G_t$  and that  $G_i$ 's are the monomer units of  $G$ . A particular case of this construction is the decomposition of a connected graph into blocks (see [9]). For more details and aspects on the polymers, we refer the reader to [1, 11, 13]. In [1] we have studied the Sombor index of polymers.

We follow the paper [1] and since we think that the similar results for elliptic Sombor index are useful for researchers, we consider the elliptic Sombor index of graphs from primary subgraphs. In Section 2, the elliptic Sombor index of some graphs are computed from their monomer units. In Section 3, we apply the results of Section 2, in order to obtain the elliptic Sombor index of families of graphs that are of importance in chemistry.


## 2 Results for Graph from Primary Subgraphs

In this section, we study the elliptic Sombor index of polymers (see [1]). By the definition of the elliptic Sombor index, we have the following easy result:


**Proposition 1.** *If  $G$  is a polymer graph with composed of monomers  $\{G_i\}_{i=1}^k$ , then*

$$ESO(G) > \sum_{i=1}^k ESO(G_i).$$

We consider some particular cases of these graphs and study their elliptic Sombor index. As an example of point-attaching graph, consider the graph  $K_m$  and  $m$  copies of  $K_n$ . Suppose that the graph  $Q(m, n)$  is obtained by identifying each vertex of  $K_m$  with a vertex of a unique  $K_n$ . The graph  $Q(5, 4)$  is shown in Figure 2. The ESO index of  $Q(m, n)$  is easy to compute.



**Figure 2.** The graph  $Q(m, n)$  and  $Q(5, 4)$ , respectively.



**Figure 3.** Link of  $n$  graphs  $G_1, G_2, \dots, G_n$

**Theorem 1.** For the graph  $Q(m, n)$  (see Figure 2), and  $n \geq 2$  we have:

$$ESO(Q(m, n)) = m((m+n-2)^2(m-1) + (n-1)^3(n-2))\sqrt{2} \\ + m(n-1)(m+2n-3)\sqrt{(m+n-2)^2 + (n-1)^2}.$$

*Proof.* There are  $\frac{m(m-1)}{2}$  edges with endpoints of degree  $m+n-2$ . Also there are  $m(n-1)$  edges with endpoints of degree  $m+n-2$  and  $n-1$  and there are  $m(n-1)(\frac{n}{2}-1)$  edges with endpoints of degree  $n-1$ . Therefore

$$ESO(Q(m, n)) = \frac{m(m-1)(2m+2n-4)}{2}\sqrt{(m+n-2)^2 + (m+n-2)^2} \\ + m(n-1)(m+2n-3)\sqrt{(m+n-2)^2 + (n-1)^2} \\ + m(n-1)(2n-2)(\frac{n}{2}-1)\sqrt{(n-1)^2 + (n-1)^2},$$

and so we have the result.  $\square$

To obtain more results, we need the following theorem.

**Theorem 2.** Suppose that  $G = (V, E)$  is a graph and  $e = uv \in E$ . If  $d_w$  is the degree of vertex  $w$  in  $G$ , then,

$$ESO(G-e) < ESO(G) - \frac{|d_u^2 - d_v^2|}{\sqrt{2}}.$$

*Proof.* First we remove edge  $e$  and find  $ESO(G-e)$ . Obviously, by adding edge  $e$  to  $G-e$  and  $(d_u + d_v)\sqrt{d_u^2 + d_v^2}$  to  $SO(G-e)$ , the  $ESO(G)$  is greater than  $ESO(G-e)$ . Since  $\sqrt{a^2 + b^2} \geq \frac{|a-b|}{\sqrt{2}}$ , so

$$ESO(G) > ESO(G-e) + (d_u + d_v)\sqrt{d_u^2 + d_v^2} \geq ESO(G-e) + \frac{(d_u + d_v)|d_u - d_v|}{\sqrt{2}},$$

and therefore we have the result.  $\square$


In the following we study the elliptic Sombor index for links of graphs, circuits of graphs, chains of graphs, and bouquets of graphs.

**Theorem 3.** Suppose that  $G$  is a polymer graph with composed of monomers  $\{G_i\}_{i=1}^k$  with respect to the vertices  $\{x_i, y_i\}_{i=1}^k$ . If  $G$  is the link of graphs (see Figure 3), then,

$$ESO(G) > \sum_{i=1}^k ESO(G_i) + \sum_{i=1}^{k-1} \frac{|d_{x_{i+1}}^2 - d_{y_i}^2|}{\sqrt{2}}.$$

*Proof.* First we remove edge  $y_1x_2$  (Figure 3). By Proposition 2, we have

$$ESO(G) > ESO(G-y_1x_2) + \frac{|d_{y_1}^2 - d_{x_2}^2|}{\sqrt{2}}.$$



**Figure 4.** Circuit of  $n$  graphs  $G_1, G_2, \dots, G_n$

If  $G'$  is the link graph related to graphs  $\{G_i\}_{i=2}^k$  with respect to the vertices  $\{x_i, y_i\}_{i=2}^k$ , then,

$$ESO(G - y_1x_2) = ESO(G_1) + ESO(G'),$$

and so,

$$ESO(G) > ESO(G_1) + ESO(G') + \frac{|d_{y_1}^2 - d_{x_2}^2|}{\sqrt{2}}.$$

By continuing this process, we have the result. □

**Theorem 4.** Let  $G_1, G_2, \dots, G_k$  be a finite sequence of pairwise disjoint connected graphs and let  $x_i \in V(G_i)$ . Suppose that  $G$  is the circuit of graphs  $\{G_i\}_{i=1}^k$  with respect to the vertices  $\{x_i\}_{i=1}^k$  and obtained by identifying the vertex  $x_i$  of the graph  $G_i$  with the  $i$ -th vertex of the cycle graph  $C_k$  (Figure 4). Then,

$$ESO(G) > \frac{|d_{x_1}^2 - d_{x_n}^2|}{\sqrt{2}} + \sum_{i=1}^k ESO(G_i) + \sum_{i=1}^{k-1} \frac{|d_{x_i}^2 - d_{x_{i+1}}^2|}{\sqrt{2}}.$$

*Proof.* First we remove edge  $x_nx_1$  (Figure 4). By Proposition 2, we have

$$ESO(G) > ESO(G - x_nx_1) + \frac{|d_{x_n}^2 - d_{x_1}^2|}{\sqrt{2}}.$$

Now we remove edge  $x_1x_2$ . So,

$$ESO(G) > ESO(G - \{x_nx_1, x_1x_2\}) + \frac{|d_{x_n}^2 - d_{x_1}^2|}{\sqrt{2}} + \frac{|d_{x_2}^2 - d_{x_1}^2|}{\sqrt{2}}.$$

Suppose that  $G'$  is the graph related to circuit graph with  $\{G_i\}_{i=2}^k$  with respect to the vertices  $\{x_i\}_{i=2}^k$  and removing the edge  $x_nx_1$ . Then we have,

$$ESO(G - \{x_nx_1, x_1x_2\}) = ESO(G_1) + ESO(G'),$$

and therefore,

$$ESO(G) > ESO(G_1) + ESO(G') + \frac{|d_{x_n}^2 - d_{x_1}^2|}{\sqrt{2}} + \frac{|d_{x_2}^2 - d_{x_1}^2|}{\sqrt{2}}.$$


By continuing this process, we have the result. □

In the following theorem we present another lower bound for the elliptic Sombor index of the circuit of graphs.

**Theorem 5.** Let  $G_1, G_2, \dots, G_k$  be a finite sequence of pairwise disjoint connected graphs and let  $x_i \in V(G_i)$ . Suppose that  $G$  is the circuit of graphs  $\{G_i\}_{i=1}^k$  with respect to the vertices  $\{x_i\}_{i=1}^k$  and obtained by identifying the vertex  $x_i$  of the graph  $G_i$  with the  $i$ -th vertex of the cycle graph  $C_k$  (Figure 4). Then,

$$ESO(G) \geq 8k\sqrt{2} + \sum_{i=1}^k ESO(G_i).$$

The equality holds if and only if for every  $1 \leq i \leq k$ ,  $G_i = K_1$ .



**Figure 5.** Chain of  $n$  graphs  $G_1, G_2, \dots, G_n$

*Proof.* Let  $d_i$  be the degree of the vertex  $x_i$  before creating  $G$ . Since  $d(x_i) = d_i + 2$ , we have:

$$\begin{aligned} ESO(G) &= (d_k + 2 + d_1 + 2) \sqrt{(d_k + 2)^2 + (d_1 + 2)^2} \\ &\quad + \sum_{i=1}^{k-1} (d_i + 2 + d_{i+1} + 2) \sqrt{(d_i + 2)^2 + (d_{i+1} + 2)^2} \\ &\quad + \sum_{i=1}^k \left( \sum_{uv \in E(G_i - x_i)} (d_u + d_v) \sqrt{d_u^2 + d_v^2} + \sum_{x_i \sim u \in G_i} (d_i + 2 + d_u) \sqrt{(d_i + 2)^2 + d_u^2} \right) \\ &\geq 4\sqrt{4+4} + \sum_{i=1}^{k-1} 4\sqrt{4+4} \\ &\quad + \sum_{i=1}^k \left( \sum_{uv \in E(G_i - x_i)} (d_u + d_v) \sqrt{d_u^2 + d_v^2} + \sum_{x_i \sim u \in G_i} (d_i + 2 + d_u) \sqrt{(d_i + 2)^2 + d_u^2} \right) \\ &= 8k\sqrt{2} + \sum_{i=1}^k SO(G_i). \end{aligned}$$

If  $G_i$  has at least one edge then the equality does not hold and so we have the result.  $\square$

**Theorem 6.** Let  $G_1, G_2, \dots, G_n$  be a finite sequence of pairwise disjoint connected graphs and let  $x_i, y_i \in V(G_i)$ . Suppose that  $C(G_1, \dots, G_n)$  is the chain of graphs  $\{G_i\}_{i=1}^n$  with respect to the vertices  $\{x_i, y_i\}_{i=1}^k$  which obtained by identifying the vertex  $y_i$  with the vertex  $x_{i+1}$  for  $i = 1, 2, \dots, n-1$  (Figure 5). Then,

(i)

$$ESO(C(G_1, \dots, G_n)) > ESO(C(G_1, \dots, G_{n-1})) + ESO(G_n - y_{n-1}) + \sum_{\substack{u \sim y_{n-1} \\ u \in V(G_n)}} \frac{|d_u^2 - d_{y_{n-1}}^2|}{\sqrt{2}}.$$

(ii)

$$ESO(C(G_1, \dots, G_n)) > ESO(C(G_1)) + \sum_{i=2}^n ESO(G_i - y_{i-1}) + \sum_{i=1}^{n-1} \sum_{\substack{u \sim y_i \\ u \in V(G_{i+1})}} \frac{|d_u^2 - d_{y_i}^2|}{\sqrt{2}}.$$

*Proof.* (i) Consider  $C(G_1, \dots, G_n)$  in Figure 5. Using inductively Theorem 2 for all edges in  $G_n$  which one of the their end vertices is  $y_{n-1}$  we have the result.

(ii) It follows by induction and Part (i).  $\square$


Similar to the Theorem 6 we have:

**Theorem 7.** Let  $G_1, G_2, \dots, G_n$  be a finite sequence of pairwise disjoint connected graphs and let  $x_i \in V(G_i)$ . Let  $B(G_1, \dots, G_n)$  be the bouquet of graphs  $\{G_i\}_{i=1}^n$  with respect to the vertices  $\{x_i\}_{i=1}^n$  and obtained by identifying the vertex  $x_i$  of the graph  $G_i$  with  $x$  (see Figure 6). Then,


$$ESO(B(G_1, \dots, G_n)) > ESO(G_1) + \sum_{i=2}^n ESO(G_i - x_i) + \sum_{i=1}^{n-1} \sum_{\substack{u \sim x_{i+1} \\ u \in V(G_{i+1})}} \frac{|d_u^2 - d_{x_{i+1}}^2|}{\sqrt{2}}.$$

### 3 Chemical Applications

In this section, using results of Section 2 to obtain the elliptic Sombor index of families of graphs that are of importance in chemistry.



**Figure 6.** Bouquet of  $n$  graphs  $G_1, G_2, \dots, G_n$  and  $x_1 = x_2 = \dots = x_n = x$



**Figure 7.** The graph  $S_{6,2,8}$

### 3.1 Spiro-Chains

Spiro-chains are defined in [7]. Using the concept of chain of graphs, a spiro-chain can be defined as a chain of cycles. We denote by  $S_{q,h,k}$  the chain of  $k$  cycles  $C_q$  in which the distance between two consecutive contact vertices is  $h$  (see  $S_{6,2,8}$  in Figure 7).

**Theorem 8.** *ESO index of the graph  $S_{q,h,k}$  for  $h \geq 2$  is:*

$$ESO(S_{q,h,k}) = (8qk - 32k + 32)\sqrt{2} + (24k - 24)\sqrt{5}.$$

*Proof.* There are  $4(k-1)$  edges with endpoints of degree 2 and 4. Also there are  $qk - 4(k-1)$  edges with endpoints of degree 2. So, we have the result.  $\square$

**Theorem 9.** *The ESO index of the graph  $S_{q,1,k}$  is:*

$$ESO(S_{q,1,k}) = (8qk + 8k - 48)\sqrt{2} + 48k\sqrt{5}.$$

*Proof.* There are  $k-2$  edges with endpoints of degree 4. Also there are  $2k$  edges with endpoints of degree 4 and 2, and there are  $qk - 3k + 2$  edges with endpoints of degree 2. Therefore we have the result.  $\square$

Cactus graphs were first known as Husimi tree, are a class of simple linear polymers. They appeared in the scientific literature some sixty years ago in papers by Husimi and Riddell [18, 20, 24]. For some aspects of parameters of cactus graphs, refer to [4, 22, 25].

As an immediate result of Theorems 8 and 9 we have the following results for cactus chains:

**Corollary 1.** (i) *If  $T_n$  is the chain triangular graph (see Figure 8) of order  $n$ , then for every  $n \geq 2$ ,  $ESO(T_n) = (32n - 48)\sqrt{2} + 32n\sqrt{5}$ .*

(ii) *If  $Q_n$  is the para-chain square cactus graph (see Figure 8) of order  $n$ , then for every  $n \geq 2$ ,  $ESO(Q_n) = 32\sqrt{2} + (48n - 48)\sqrt{5}$ .*


(iii) *If  $O_n$  is the para-chain square cactus (see Figure 9) graph of order  $n$ , then for every  $n \geq 2$ ,  $ESO(O_n) = (40n - 48)\sqrt{2} + 24n\sqrt{5}$ .*

(iv) *If  $O_n^h$  is the Ortho-chain graph (see Figure 9) of order  $n$ , then for every  $n \geq 2$ ,  $ESO(O_n^h) = (56n - 48)\sqrt{2} + 24n\sqrt{5}$ .*

(v) *If  $L_n$  is the para-chain hexagonal cactus graph (see Figure 10) of order  $n$ , then for every  $n \geq 2$ ,  $ESO(L_n) = (16n + 32)\sqrt{2} + (48n - 48)\sqrt{5}$ .*



**Figure 8.** Chain triangular cactus  $T_n$  and para-chain square cactus  $Q_n$



**Figure 9.** Para-chain square cactus  $O_n$  and ortho-chain graph  $O_n^h$

(vi) If  $M_n$  is the Meta-chain hexagonal cactus graph (see Figure 10) of order  $n$ , then for every  $n \geq 2$ ,  $ESO(M_n) = (16n + 32)\sqrt{2} + (48n - 48)\sqrt{5}$ .

### 3.2 Polyphenylenes

Similar to the definition of the spiro-chain  $S_{q,h,k}$ , we can define the graph  $L_{q,h,k}$  as the link of  $k$  cycles  $C_q$  in which the distance between the two contact vertices in the same cycle is  $h$  (see  $L_{6,2,4}$  in Figure 11).

**Theorem 10.** The ESO index of the graph  $L_{q,h,k}$ , for  $h \geq 2$  is:

$$ESO(L_{q,h,k}) = (8qk - 14k + 14)\sqrt{2} + (20k - 20)\sqrt{13}.$$

*Proof.* There are  $k - 1$  edges with endpoints of degree 3. Also there are  $4(k - 1)$  edges with endpoints of degree 3 and 2, and there are  $qk - 4(k - 1)$  edges with endpoints of degree 2. Therefore we have the result.  $\square$


**Theorem 11.** The ESO index of the graph  $L_{q,1,k}$  is:

$$ESO(L_{q,1,k}) = (8qk + 12k - 38)\sqrt{2} + 10k\sqrt{13}.$$

*Proof.* There are  $2k - 3$  edges with endpoints of degree 3. Also there are  $2k$  edges with endpoints of degree 3 and 2, and there are  $qk - 3k + 2$  edges with endpoints of degree 2. Therefore we have the result.  $\square$

### 3.3 Triangulanes

We want to obtain the elliptic Sombor index of the triangulane  $T_k$  defined pictorially in [21]. The triangulane  $T_k$  is defined recursively in a manner that is useful in our approach. First define recursively an auxiliary family of triangulanes  $G_k$  ( $k \geq 1$ ). Let  $G_1$  be a triangle and denote one of its vertices by  $y_1$ . Define  $G_k$  ( $k \geq 2$ ) as the circuit of the graphs  $G_{k-1}$ ,  $G_{k-1}$ , and  $K_1$  and denote by  $y_k$  the vertex where  $K_1$  has been placed. The graphs  $G_1$ ,  $G_2$  and  $G_3$  are shown in Figure 12.



**Figure 10.** Para-chain  $L_n$  and Meta-chain  $M_n$




Figure 11. The graph  $L_{6,2,4}$




Figure 12. Graphs  $G_1$ ,  $G_2$  and  $G_3$





Figure 13. Graphs  $T_3$



**Figure 14.** Graphs  $F$  and  $G_1$ , respectively.



**Figure 15.** Graphs  $G_2$  and  $G_3$ , respectively.

**Theorem 12.** The ESO index of the graph  $T_k$  (see  $T_3$  in Figure 13) is:

$$ESO(T_k) = (288(2^{k-1} - 1) + 24(2^{k-1}) + 96)\sqrt{2} + 36(2^k)\sqrt{5}.$$

*Proof.* By recursive structure of the graph  $T_k$ , observe that there are  $3 + 3 \sum_{n=0}^{k-2} 3(2^n)$  edges with endpoints of degree 4. Also there are  $3(2^k)$  edges with endpoints of degree 4 and 2, and there are  $3(2^{k-1})$  edges with endpoints of degree 2. Therefore

$$ESO(T_k) = (3 + 9 \sum_{n=0}^{k-2} 2^n)(8)\sqrt{16+16} + 3(2^k)(6)\sqrt{16+4} + 3(2^{k-1})(4)\sqrt{4+4},$$

and so we have the result.  $\square$

### 3.4 Nanostar Dendrimers

In this subsection, we want to compute the elliptic Sombor index of the nanostar dendrimer  $D_k$  defined in [3]. In order to define  $D_k$ , we follow [9]. First we define recursively an auxiliary family of rooted dendrimers  $G_k$  ( $k \geq 1$ ). We need a fixed graph  $F$  defined in Figure 14, we consider one of its endpoint to be the root of  $F$ .


The graph  $G_1$  is defined in Figure 14, the leaf being its root. Now we define  $G_k$  ( $k \geq 2$ ) the bouquet of the following three graphs:  $G_{k-1}$ ,  $G_{k-1}$ , and  $F$  with respect to their roots; the root of  $G_k$  is taken to be its unique leaf (see  $G_2$  and  $G_3$  in Figure 15). Finally, we define  $D_k$  ( $k \geq 1$ ) as the bouquet of three copies of  $G_k$  with respect to their roots ( $D_2$  is shown in Figure 16, where the circles represent hexagons).

**Theorem 13.** The ESO index of the dendrimer  $D_3[n]$  (see  $D_3[2]$  in Figure 16) is:

$$ESO(D_3[n]) = (468 \times 2^n + 204)\sqrt{2} + (90 \times 2^n + 30)\sqrt{13}.$$

*Proof.* There are  $3 + 9 \sum_{k=0}^{n-1} 2^k$  edges with endpoints of degree 3. Also there are  $6 + 18 \sum_{k=0}^{n-1} 2^k$  edges with endpoints of degree 3 and 2, and there are  $12 + 18 \sum_{k=0}^{n-1} 2^k$  edges with endpoints of degree 2. Therefore

$$ESO(D_3[n]) = (3 + 9 \sum_{k=0}^{n-1} 2^k)(6)\sqrt{9+9} + (6 + 18 \sum_{k=0}^{n-1} 2^k)(5)\sqrt{9+4}$$



**Figure 16.** Nanostar  $D_2$  and  $D_3[2]$ , respectively.

$$+ (12 + 18 \sum_{k=0}^{n-1} 2^k)(4)\sqrt{4+4},$$

and we have the result. □

## Authors' Contributions

All authors have the same contribution.

## Data Availability

The manuscript has no associated data or the data will not be deposited.

## Conflicts of Interest

The authors declare that there is no conflict of interest.

## Ethical Considerations

The authors have diligently addressed ethical concerns, such as informed consent, plagiarism, data fabrication, misconduct, falsification, double publication, redundancy, submission, and other related matters.

## Funding

This research did not receive any grant from funding agencies in the public, commercial, or nonprofit sectors.

## References

- [1] S. Alikhani, N. Ghanbari, Sombor index of polymers, *MATCH Commun. Math. Comput. Chem.*, 86, 715–728 (2021).
- [2] S. Alikhani, N. Ghanbari, M. A. Dehghanizadeh, Elliptic Sombor energy of a graph, submitted. Available at <https://arxiv.org/abs/2404.18622>.
- [3] S. Alikhani, M. A. Iranmanesh, Hosoya polynomial of dendrimer nanostar  $D_3[n]$  *MATCH Commun. Math. Comput. Chem.* 71, 395–405 (2014).
- [4] M. Chellali, Bounds on the 2-domination number in cactus graphs, *Opuscula Math.* 2, 5–12 (2006).

- [5] K. C. Das, A. S. Cevik, I. N. Cangul, Y. Shang, On Sombor index, *Symmetry* 13, 140 (2021).
- [6] H. Deng, Z. Tang, R. Wu, Molecular trees with extremal values of Sombor indices, *Int. J. Quantum Chem.*, 121, e26622 (2021).
- [7] M. V. Diudea, I. Gutman, J. Lorentz, *Molecular Topology*, Nova Science Publishers Huntington, NY, USA, (2001).
- [8] R. Cruz, I. Gutman, J. Rada, Sombor index of chemical graphs, *Appl. Math. Comp.*, 399, 126018 (2021).
- [9] D. Emeric, S. Klavžar, Computing Hosoya polynomials of graphs from primary subgraphs, *MATCH Commun. Math. Comput. Chem.*, 70, 627–644 (2013).
- [10] C. Espinal, I. Gutman, J. Rada, Elliptic Sombor index of chemical graphs, *Commun. Comb. Optim.*, 1-11 (2024).
- [11] N. Ghanbari, On the Graovac-Ghorbani and atom-bond connectivity indices of graphs from primary subgraphs, *Iran. J. Math. Chem.*, 13(1), 45–72 (2022).
- [12] N. Ghanbari, On the Sombor characteristic polynomial and Sombor energy of a graph, *Comp. Appl. Math.*, 41, 242 (2022).
- [13] N. Ghanbari, S. Alikhani, Mostar index and edge Mostar index of polymers, *Comp. Appl. Math.*, 40, 260 (2021).
- [14] N. Ghanbari, S. Alikhani, Sombor index of certain graphs, *Iranian J. Math. Chem.*, 12(1), 27–37 (2021).
- [15] N. Ghanbari, S. Alikhani, Sombor polynomial (function) of a graph, *J. Discrete Math. Sci. Cryptogr.*, 27(3), 983–998 (2024).
- [16] I. Gutman, Geometric approach to degree based topological indices, *MATCH Commun. Math. Comput. Chem.* 86(1) 11-16 (2021).
- [17] I. Gutman, B. Furtula, M. Sinan Oz, Geometric approach to vertex-degree-based topological indices Elliptic Sombor index, theory and application, *Quantum Chem.* 124 (2) e27346 (2024).
- [18] F. Harary, B. Uhlenbeck, On the number of Husimi trees, I, *Proc. Nat. Acad. Sci.* 39 315-322 (1953).
- [19] S. Hayat, M. Arshad, A. Khan, Graphs with given connectivity and their minimum Sombor index having applications to QSPR studies of monocarboxylic acids, *Heliyon* 10 e23392 (2024).
- [20] K. Husimi, Note on Mayer's theory of cluster integrals, *J. Chem. Phys.* 18 682-684 (1950).
- [21] M.H. Khalifeh, H. YousefiAzari, A.R. Ashrafi, Computing Wiener and Kirchhoff indices of a triangulane, *Indian J. Chem.* 47A 1503-1507 (2008).
- [22] S. Majstorović, T. Došlić, A. Klobučar,  $k$ -domination on hexagonal cactus chains, *Kragujevac J. Math.*, 36, No 2, 335-347 (2012).
- [23] I. Redžepović, Chemical applicability of Sombor indices, *J. Serb. Chem. Soc.* 86, 445-457 (2021).
- [24] R.J. Riddell, Contributions to the theory of condensation, Ph.D. Thesis, Univ. of Michigan, Ann Arbor, 1951.
- [25] A. Sadeghieh, N. Ghanbari, S. Alikhani, Computation of Gutman index of some cactus chains, *Elect. J. Graph Theory Appl.* 6(1) 138-151 (2018).
- [26] Z. Wang, Y. Mao, Y. Li, B. Furtula, On relations between Sombor and other degree-based indices, *J. Appl. Math. Comput.* 68, 1-17 (2022).
- [27] H. Wiener, Structural determination of the Paraffin Boiling Points, *J. Amer. Chem. Soc.*, 69, 17-20 (1947).